

Optical Properties, devices ...

Ram Seshadri MRL 2031, x6129, seshadri@mrl.ucsb.edu

Some basics:

Electromagnetic radiation has electric and magnetic field components that are perpendicular to each other and to the direction of propagation. The propagation velocity is c , and this is constant for all electromagnetic radiation in vacuum: $c = 2.997\,924\,58 \times 10^8$ m/s.

c is related to the vacuum permittivity ϵ_0 and the vacuum permeability μ_0 in the following manner:

$$c = \frac{1}{\sqrt{\epsilon_0 \mu_0}}$$

The values of the permeability and the permittivity are:

$$\epsilon_0 = 8.854\,187\,817 \times 10^{-12} \text{ F m}$$

$$\mu_0 = 4\pi \times 10^{-7} \text{ N A}^{-2}$$

c is equal to the product of the frequency and the wavelength:

$$c = \nu \lambda$$

The energy of any light photon is proportional to the frequency of the photon so:

$$E = h\nu$$

where ν is in Hz (s^{-1}) and h is the Planck constant $h = 6.63 \times 10^{-34}$ Js. It is also possible then to write:

$$E = \frac{hc}{\lambda}$$

Electromagnetic radiation with different frequencies have correspondingly, different energies and wavelengths. Visible electromagnetic radiation (“light”) has wavelengths ranging from 400 to 700 nm.

Light and metals — reflection:

Since metals have free electrons, all energies are absorbed and re-emitted. Hence the reflectivity. Yellow/red metals such as Cu and Au are so colored because they absorb a little in the blue.

Refraction:

The refractive index n of a material is the ratio of the velocity of light in vacuum (always c) to the velocity of light in that material v :

$$n = \frac{c}{v}$$

If the material has a dielectric constant ϵ and permeability μ , then

$$v = \frac{1}{\sqrt{\epsilon\mu}}$$

Therefore:

$$n = \frac{\sqrt{\epsilon\mu}}{\sqrt{\epsilon_0\mu_0}} = \sqrt{\epsilon_r\mu_r}$$

Since for most substances, $\mu_r \sim 1$,

$$n \sim \sqrt{\epsilon_r}$$

Even transparent materials reflect some portion of the light that falls on them. If the light falls normally from a medium of refractive index n_1 to the medium of refractive index n_2 , then the proportion of light reflected is


$$R = \left(\frac{n_2 - n_1}{n_2 + n_1} \right)^2$$

Color:

The color of a material is due to the light that is reflected by it. If a material absorbs at one end of the spectrum, its color is usually the other end. There are (usually) three processes by which a material may be colored: Absorption with atomic or defect levels, absorption across the band gap, and due to patterning (interference or diffraction as in a soap film or a peacock's feather).

The creation of light:

A lot of light is created using heat (black-body radiation). In rare cases, chemical changes are directly converted to light (fireflies) and in less rare cases, electrical energy is converted to light in a light emitting diode:



Note that if the device is turned on its head, one has a solar cell.

Lasers: These rely on the concepts of stimulated emission and population inversion.