

MATRL 218: Assignment 6

Ram Seshadri (seshadri@mrl.ucsb.edu)

1. Explain how you would use a *local* bonding picture to sketch the density of states of Si (filled valence band and empty conduction band) by recognizing that with 4 electrons per Si and 4 bonds going out of every Si, the compound is electron precise.
2. Can you extend this analysis to amorphous Si?
3. How should one think of the electronic structure of SiO₂, which has an enormous band gap? Remember that you need to think of the separation between the centers of states that form the valence and conduction band (that can usually be attributed to electronegativity differences) *and* you need to think of the width of the VB and CB (which arises from dispersion — the orbitals involved and the nature of the structure).
4. CaF₂ has an even larger band gap than SiO₂, which is of great consequence in high-end camera lenses (larger band gaps mean reduced chromatic aberration). Suggest using ideas of electronegativity and band width, why this might be so.
5. I-III-VI semiconductors such as CuInSe₂, II-VI semiconductors such as CdSe, and III-V semiconductors such as GaAs all have the same rules of electron precision as Si in the diamond structure. Explain these.
6. The compound LaCoO₃ is a diamagnetic semiconductor at low temperatures, but when heated, displays metallic behavior and some magnetism. In contrast, the compound LaRhO₃ is always insulating. Use crystal field arguments for d^6 Co³⁺ and Rh³⁺ to explain this.